

SUI-64-36

2 Copies

UNPUBLISHED PRELIMINARY DATA

The Alfvén Invariant in the
Field of a Magnetic Unipole

J. A. Van Allen
Department of Physics and Astronomy
University of Iowa, Iowa City

FACILITY FORM 602

N 65 14958
(ACCESSION NUMBER)
4
(PAGES)
CR-60141
(NASA CR OR TMX OR AD NUMBER)

(THRU)
1
(CODE)
23
(CATEGORY)

GPO PRICE \$ _____

OTS PRICE(S) \$ _____

Hard copy (HC) \$ 1.00

Microfiche (MF) \$ 0.50

November 1964

The Alfvén Invariant in the
Field of a Magnetic Unipole

J. A. Van Allen
Department of Physics and Astronomy
University of Iowa, Iowa City

It is well known [Alfvén, 1950] [Northrop, 1963] that an isolated, electrically-charged particle moves in a static magnetic field in such a way that the quantity

$$\frac{p_{\perp}^2}{2m_0 B} \quad (1)$$

is an adiabatic invariant of the motion provided that

$$\rho \left| \frac{\text{grad } B}{B} \right| \ll 1. \quad (2)$$

In (1) and (2), p_{\perp} is the component of the particle's longitudinal, relativistic momentum p perpendicular to the magnetic vector B , m_0 is its rest mass, and ρ is the radius of curvature of the osculating circle to its trajectory. Since $p_{\perp} = p \sin \alpha$, where α is the angle between p and B (instantaneous pitch angle), and since scalar p and m_0 are constants of the motion, it follows from the adiabatic invariance of quantity (1) that

$$\frac{\sin^2 \alpha}{B} = \text{Adiabatic Constant.} \quad (3)$$

The purpose of the present note is to call attention to the fact that $\sin^2 \alpha/B$ is a rigorous constant of the motion of an isolated, electrically-charged particle in the field of a magnetic unipole.

It was shown by Darboux [1878] that the equilibrium form of a flexible, inextensible, current-carrying wire which is under tension and in the field of a magnetic unipole is a geodesic on a circular cone, whose apex is at the unipole. Later, Poincare [1896] pointed out that the differential equations of the trajectory of an electrically-charged particle are of identical form to those of Darboux's current-carrying wire provided one sets the ratio of the tension to the current equal to the magnetic rigidity of the particle. Hence, the trajectory of an electrically-charged particle is also a geodesic on a circular cone whose apex is at the unipole. The complete properties of the trajectory are determined by exact integration of the equations of motion with given initial conditions. A simple, elegant discussion using vector notation is given by Ferraro [1956].

It follows from the properties of geodesics on circular cones [Struik, 1961] that

$$r \sin \theta \sin \alpha = \text{constant} \quad (4)$$

where r is the radial distance from the apex of the cone to a point on the geodesic, θ is the half angle of the cone, and α is the angle between the geodesic line and the element of the cone through the point. For a unipole

$$B r^2 = \text{constant} \quad (5)$$

and B lies along the element of the cone.

From (4) and (5) it is seen that the Alfven quantity $\sin^2 \alpha / B$ is a rigorous constant of the motion. Q.E.D.

References

- Alfven, H., "Cosmical Electrodynamics", Oxford at the Clarendon Press, 1950.
- Darboux, G., "Probleme de Mecanique", Bulletin des Sciences Mathematiques et Astronomiques Ser. 2, Vol. 2, 433-436, 1878.
- Ferraro, V. C. A., "Electromagnetic Theory", Athlone Press, University of London, 1956.
- Northrop, T. G., "The Adiabatic Motion of Charged Particles" Interscience Publishers, New York, 1963.
- Poincare, H., Remarques sur une experience de M. Birkeland, Comptes Rendus, Acad. Sci. Paris 123, 530-533, 1896.
- Struik, D. J., "Differential Geometry", Addison-Wesley Publishing Company, Inc., Reading, Mass., 2nd Ed., 1961.